

兔抗 MAP3K8(Ab-400) 多克隆抗体

- 中文名称：兔抗 MAP3K8(Ab-400) 多克隆抗体
- 英文名称：Anti-MAP3K8(Ab-400) rabbit polyclonal antibody
- 别名：COT; EST; ESTF; TPL2; MEKK8; Tpl-2; c-COT
- 相关类别：一抗
- 储存：冷冻（-20℃）避光
- 宿主：Rabbit
- 抗原：MAP3K8(Ab-400)
- 反应种属：Human
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:	This gene is an oncogene that encodes a member of the serine/threonine protein kinase family. The encoded protein localizes to the cytoplasm and can activate both the MAP kinase and JNK kinase pathways. This protein was shown to activate I κ B kinases, and thus induce the nuclear production of NF- κ B. This protein was also found to promote the production of TNF- α and IL-2 during T lymphocyte activation. This gene may also utilize a downstream in-frame translation start codon, and thus produce an isoform containing a shorter N-terminus.
Applications:	WB, IHC
Name of antibody:	MAP3K8(Ab-400)

Immunogen:	Synthesized non-phosphopeptide derived from human MA P3K8 around the phosphorylation site of serine 400 (C-Q-S(p)-L-D).
Full name:	mitogen-activated protein kinase kinase kinase 8
Synonyms :	COT; EST; ESTF; TPL2; MEKK8; Tpl-2; c-COT
SwissProt:	P41279
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	53 kDa
WB Positive control:	COLO cells treated with Insulin (0.01U/ml, 15mins) lysate
WB Recommended dilution:	500-3000



