

兔抗 STAP1 多克隆抗体

中文名称: 兔抗 STAP1 多克隆抗体

英文名称: Anti-STAP1 rabbit polyclonal antibody

别名: BRDG1; STAP-1

抗原: STAP1

储存: 冷冻 (-20℃)

宿主: Rabbit

反应种属: Human, Mouse

相关类别: 一抗

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	The protein encoded by this gene contains a proline-rich region, a pleckstrin homology (PH) domain, and a region in the carboxy terminal half with similarity to the Src Homology 2 (SH2) domain. This protein is a substrate of tyrosine-protein kinase Tec, and its interaction with tyrosine-protein kinase Tec is phosphorylation-dependent. This protein is thought to participate in a positive feedback loop by upregulating the activity of tyrosine-protein kinase Tec. Variants of this gene have been associated with autosomal-dominant hypercholesterolemia (ADH), which is characterized by elevated low-density lipoprotein cholesterol levels and in increased risk of coronary vascular disease. Alternative splicing results in multiple transcript variants.
Applications:	ELISA, IHC

Name of antibody:	STAP1
Immunogen:	Fusion protein of human STAP1
Full name:	signal transducing adaptor family member 1
Synonyms:	BRDG1; STAP-1
SwissProt:	Q9ULZ2
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer and human esophagus cancer
IHC Recommend dilution:	25-100



