

## H1-6 抗原(重组蛋白)

- 中文名称: H1-6 抗原 (重组蛋白)
- 英文名称: H1-6 Antigen (Recombinant Protein)

别 名: H1.6 linker histone, cluster member; H1t; H1.6; H1FT; HIST1H1T; dJ221C16.2

- 相关类别: 抗原
- 储 存: 冷冻(-20℃)

## 概述

Fusion protein corresponding to a region derived from 1-207 amino acids of human H1-6

## 技术规格

<b>F</b> II	111 C lialaan bistore shusten meenkan
Full name:	H1.6 linker histone, cluster member
Synonyms:	H1t; H1.6; H1FT; HIST1H1T; dJ221C16.2
Swissprot:	P22492
Gene Accession:	BC130350
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	Histones are basic nuclear proteins responsible for nucleosome struc ture of the chromosomal fiber in eukaryotes. Two molecules of eac h of the four core histones (H2A, H2B, H3, and H4) form an octam er, around which approximately 146 bp of DNA is wrapped in repea ting units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless an d encodes a replication-dependent histone that is a member of the histone H1 family. Transcripts from this gene lack polyA tails but in stead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by R



efSeq, Aug 2015]